



1. Consider the two reduction processes and their standard reduction potentials (E°).



A) Write the net ionic equation for a Galvanic/voltaic cell based on these reactions.


B) Determine the value of the E°_{cell} .

C) Determine the value of the standard free energy change of the cell ($\Delta G^\circ_{\text{cell}}$).

D) Determine the equilibrium constant (K) for the reaction.


E) Given below is an unlabeled diagram. Label the following components in the diagram:

- The solid electrodes on the anode and cathode sides.
- The ions in solutions on the anode and cathode sides.
- The direction of the flow of electrons through the voltmeter and wire.
- The direction of the flow of cations and anions in a salt bridge made of $\text{KNO}_3 (\text{aq})$.



F) Write the cell diagram for this electrochemical cell.

2. You have constructed a Galvanic cell with the following reaction under standard conditions.





What will the potential of the cell be when 0.50 M of $\text{Cu}^{2+} (\text{aq})$ has reacted?

Assume that volume and temperature do not change.

3. Consider an electrochemical cell with the following cell diagram at 298.15 K.



Given the following E° values, determine whether each statement is true or false.



- A) E_{cell} is a smaller value than E°_{cell} .
- B) The oxidation reaction takes place at the anode.
- C) Adding 1.0 L of water to both the anodic and cathodic solutions will increase the cell potential.
- D) Decreasing the concentration of Ni^{2+} will increase the cell potential.
- E) Increasing the concentration of Cu^{2+} will increase the cell potential.
- F) Using a Pt electrode in place of the Ni electrode will not change the cell potential.
- G) The mass of the Cu electrode will decrease over time.